



US006214875B1

(12) United States Patent
Yang

(10) Patent No.: US 6,214,875 B1

(45) Date of Patent: Apr. 10, 2001

**(54) ANTICANCER EFFECTS OF SPECIFIC
BRANCHED-CHAIN FATTY ACIDS AND
RELATED PRODUCTION PROCESS**

(76) Inventor: Zhenhua Yang, 3008 Andalucia Dr.,
West Covina, CA (US) 91791

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: 09/173,681

(22) Filed: Oct. 16, 1998

Related U.S. Application Data

(60) Provisional application No. 60/081,712, filed on Apr. 14,
1998.

(51) Int. Cl.: A61K 31/20

(52) U.S. Cl.: 514/558; 514/2; 514/8;
514/557; 514/560; 426/61; 424/115; 424/116;
424/123

(58) Field of Search: 514/2, 8, 558,
514/560, 557; 426/61; 424/115, 116, 123

(56) References Cited

U.S. PATENT DOCUMENTS

4,877,739 10/1989 Yang .
4,985,466 1/1991 Deguchi .

FOREIGN PATENT DOCUMENTS

2020633 8/1997 (CA) .

OTHER PUBLICATIONS

Chemical Abstracts 118:139830, "Higher Fatty Acids as
Anticancer Agents and Anticancer Agent Enhancers", Aug.
1995.*

Hansen, et al., "The Branched-chain Fatty Acids of Mutton
Fat. The Isolation of (+)-12-methyltetradecanoic Acid and
of 13-methyltetradecanoic Acid," *Biochem. J.*, vol. 53, pp.
374-378 (1953).

Kaneda, T., "Fatty Acids of the Genus *Bacillus*: an Example
of Branched-Chain Preference," *Bacteriological Reviews*,
vol. 41, No. 2, pp. 391-418, (1977).

Hansen, R.P., Shorland, F.B. and Cooke, N.J. The Branched-
Chain Fatty Acids of Mutton Fat, 2. The Isolation of
(+)-12-Methyltetradecanoic Acid and of 13-Methyltetra-
decanoic Acid. *Biochem. J.* 53:374 (1953).

Hansen R.P., Shorland, F.B. and Cooke, N.J. The Branched-
Chain Fatty Acids of Butterfat, 4. The Isolation of
(+)-12-Methyltetradecanoic Acid and of 13-Methyltetra-
decanoic Acid. *Biochem. J.* 57:297 (1954).

Klein, R.A., Halliday, D. and Pittet, P.G. The Use of
13-Methyltetradecanoic Acid as an Indicator of Adipose
Tissue Turnover. *Lipids* 15:572 (1980).

Pittet, P.G., Bessart, T., Jequier, E., Philippoussian, G. and
Liardon, R. Adipose Tissue Labelling in Man, Using a
Structurally-Labelled Fatty Acid as a Tracer. *Intern. J. Vit.
Nutr. Res.* 53:115 (1982).

Faung, S.T., Chiu, L. and Wang C.T., Platelet Lysis and
Functional Perturbation by 13-Methyl Myristate, the Major
Fatty Acid in *Flavobacterium Ranacida*. *Thromb. Res.* 81:91
(1996).

Wells, J.M., Butterfield, J.E. and Reveal, L.G., Identification
of Bacteria Associated with Postharvest Diseases of Fruits
and Vegetables by Cellular Fatty Acid Composition: An
Expert System for Personal Computers. *Phytopathology*
83:445 (1993).

Dee, S.B., Karr, D.E., Hollis, D. and Woss, C.W. Cellular
Fatty Acids of *Capnocytophaga* Species. *J. Clin. Microbiol.*
16:779 (1982).

Heerdt, B.G., Houston, M.A. and Augenlicht, L.H. Potentiation
by Specific Short-Chain Fatty Acids of Differentiation
and Apoptosis in Human Carcinoma Cell Lines. *Cancer Res.*
54:3288 (1994).

* cited by examiner

Primary Examiner—Frederick Krass

(74) Attorney, Agent, or Firm—Oblon, Spivak, McClelland,
Maier & Neustadt, P.C.

(57) ABSTRACT

A group of specific branched-chain fatty acids, with significant
anticancer effects on human and animals; methods of
making using either chemical synthesis or biosynthesis
methods; and methods of treating cancer.

23 Claims, 7 Drawing Sheets

(3 of 7 Drawing Sheet(s) Filed in Color)